

PHASE ANGLE BIBLIOGRAPHY



BASIC SCIENCE

- 1) Lukaski, HC; Biological Indexes Considered in the Derivation of the Bioelectrical Impedance Analysis. American Journal of Clinical Nutrition; 1996; 64(supplement); page 397s – 404s
- 2) Foster, KR & Lukaski, HC; Whole-body Impedance-what does it measure? American Journal of Clinical Nutrition; 1996; 64(supplement); page 388s – 396s
- 3) Cole, KS; Electrical Phase Angle of Cell Membranes. The Journal of General Physiology, July 20, 1932, p 641 – 649
- 4) Hoffer, EC, et al; Correlation of Whole-body impedance with Total Body Water Volume. Journal of Applied Physiology, Volume 27, Number 4, October 1969, p 531 – 534.
- 5) Kyle UG, et al; Bioelectrical Impedance Analysis—Part I: Review of Principles and Methods Clinical Nutrition 2004; 23:1226–1243
- 6) Kyle UG, et al; Bioelectrical Impedance Analysis—Part II: Utilization in Clinical Practice Clinical Nutrition 2004; 23:1430–1453
- 7) Barbosa-Silva, MCG, et al; Bioelectrical Impedance Analysis: Population Reference Values for Phase Angle by Age and Sex. American Journal of Clinical Nutrition; 2005;82:49–52
- 8) Kyle UG, et al; Percentiles (10, 25, 75 and 90th) for Phase Angle (PhA), Determined by Bioelectrical Impedance (BIA), in 2740 Healthy Adults Aged 20–75 yr. Clinical Nutrition 2004; 23:758
- 9) Chumlea, WC, et al; Body Composition Estimates from NHANES III Bioelectrical Impedance Data. International Journal of Obesity (2002) 26, 1596–1609
- 10) Bosy-Westphal, A, et al; Phase Angle From Bioelectrical Impedance Analysis: Population Reference Values by Age, Sex, and Body Mass Index. JPEN, Jul-Aug 2006

ONCOLOGY

- 1) Gupta, D, et al; Bioelectrical Impedance Phase Angle as a Prognostic Indicator in Advanced Pancreatic Cancer. British Journal of Nutrition (2004), 92, 957–962
- 2) Toso S, Piccoli A, et al; Altered Tissue Electric Properties in Lung Cancer Patients as Detected by Bioelectric Impedance Vector Analysis. Nutrition 2000; 16:120–4

- 3) De Luis, D.A, et al; Tissue Electric Properties in Head and Neck Cancer Patients' Annals of Nutrition and Metabolism 2006; 50:7-10 (DOI: 10.1159/000089484)
- 4) Toso S., Piccoli A, et al; Bioimpedance Vector Pattern in Cancer Patients Without Disease Versus Locally Advanced or Disseminated Disease. Nutrition 2003; 19: 510–4
- 5) Gupta, D. et al; Bioelectrical Impedance Phase Angle in Clinical Practice: Implications for Prognosis in Advanced Colorectal Cancer. American Journal of Clinical Nutrition; 2004; 80:1634–8
- 6) Gupta, D. et al; Bioelectrical Impedance Phase Angle as a Prognostic Indicator in Breast Cancer. BMC Cancer 2008, 8:249
- 7) Gupta, D. et al; Bioelectrical Impedance Phase Angle in Clinical Practice: Implications for Prognosis in Stage IIIB and IV Non-Small Cell Lung Cancer. BMC Cancer 2009, 9:37
- 8) Davis, MP. et al; Bioelectric Impedance Phase Angle Changes During Hydration and Prognosis in Advanced Cancer, American Journal of Hospital and Palliative Care, 2009 Jan 30. (Electronic publication ahead of print)
- 9) Davis, MP., et al; Phase Angle is a Better Predictor of Survival Than Muscle Mass in Advanced Cancer, MASCC/ISOO 2009 International Symposium, Supportive Care in Cancer, Abstract Number: 270, Last Modified: February 13 2009
- 10) Santarpia, L., et al; Prognostic Significance of Bioelectrical Impedance Phase Angle in Advanced Cancer: Preliminary Observations, Nutrition 25 (2009) 930–931
- 11) Paiva, SI. et al; Standardized Phase Angle from BIA as Prognostic Factor for Survival in Patients with Cancer, Support Care Cancer, DOI 10.1007/s00520-009-0798-9, Received: 27 July 2009 / Accepted: 8 December 2009, Springer-Verlag
- 12) Pelzer, U., et al; Parenteral Nutrition Support for Patients with Pancreatic Cancer. Results of a Phase II Study, BMC Cancer 2010, 10:86

PROGNOSIS

- 1) Mushnick, R. et al; Relationship of Bioelectrical Impedance Parameters to Nutrition and Survival in Peritoneal Dialysis Patients. Kidney International (2003)83, Suppl., S53–S56
- 2) Schwenk, A. et al; Bioelectrical Impedance Analysis Predicts Outcome in Patients with Suspected Bacteremia. Infection (1998)26, 277–282
- 3) Shime, N. et al; Bioelectrical Impedance Analysis for Assessment of Severity of Illness in Pediatric Patients After Heart Surgery. Critical Care Med 2002 Vol. 30, No. 3
- 4) Scheltinga M.R., et al; Altered Cell Membrane Function in Critical Illness can be Characterized by Measuring Body Reactance, Surgery Forum 1990; 41:43–44

- 5) Swaraj, S. et al; Bioelectrical Impedance Analysis as a Predictor for Survival in Patients with Systemic Inflammatory Response Syndrome. Critical Care 2003, 7 (Supplement2):P185
- 6) Paterna, S. et al; Changes in Brain Natriuretic Peptide Levels and Bioelectrical Impedance Measurements After Treatment With High-Dose Furosemide and Hypertonic Saline Solution Versus High-Dose Furosemide Alone in Refractory Congestive Heart Failure A Double-Blind Study. Journal of the American College of Cardiology, Vol. 45, No. 12, 2005 :1997–2003
- 7) Frankenfield, DC, et al; Bioelectrical Impedance Plethysmographic Analysis of Body Composition in Critically Injured and Healthy Subjects. American Journal of Clinical Nutrition 1999; 69:426–31
- 8) Scanferla, F, et al; On-Line Bioelectric Impedance During Haemodialysis: Monitoring of Body Fluids and Cell Membrane Status. Nephrology Dialysis & Transplant Supplement, 1 (1990) 167-170
- 9) Pupim, LB, et al; Uremic Malnutrition as a Predictor of Death Independent of Inflammatory Status, Kidney International, Volume 66, 2004, p 2054-2060
- 10) Nagano, M, et al; The Validity of Bioelectrical Impedance Phase Angle for Nutritional Assessment in Children. Journal of Pediatric Surgery, Volume 35, No 7 (July), 2000: pp 1035-1039
- 11) Maggiore, Q, et al; Nutritional and Prognostic Correlates of Bioimpedance Indexes in Hemodialysis Patients, Kidney International, 50:2103–2108, 1996
- 12) Chertow GM, et al; Phase Angle Predicts Survival in Hemodialysis Patients. Journal of Renal Nutrition 7:204–207, 1997
- 13) Pillon, L, et al; Vector Length as a Proxy for the Adequacy of Ultrafiltration in Hemodialysis. Kidney International, Vol. 66 (2004), pp. 1266–1271
- 14) Di Iorio, B R, et al; A Systematic Evaluation of Bioelectrical Impedance Measurement after Hemodialysis Session. Kidney International, Vol. 65 (2004), pp. 2435–2440
- 15) Di Iorio, B R, et al; Charlson Comorbidity Index is a Predictor of Outcomes in Incident Hemodialysis Patients and Correlates with Phase Angle and Hospitalization. International Journal of Artificial Organs 2004 Apr; 27 (4):330-6
- 16) Desport, JC, et al; Phase Angle is a Prognostic Factor for Survival in Amyotrophic Lateral Sclerosis. Amyotrophic Lateral Sclerosis 2008 Oct; 9(5):273-8
- 17) David, S., et al; Prospective Evaluation of an In-Centre Conversion from Conventional Hemodialysis to an Intensified Nocturnal Strategy. Nephrology, Dialysis and Transplantation, 2009 July 24 (7) 2232-40
- 18) Guida, B. et al; Abnormalities of Bioimpedance Measures in Overweight and Obese Hemodialyzed Patients. International Journal of Obesity (2001) 25, 265-72

- 19) Silva-Tinoco, R, et al; Developing Thyroid Disorders is Associated with Poor Prognosis Factors in Patients with Stable Chronic Heart Failure, International Journal of Cardiology, 2009 February, (E-Pub)
- 20) Cinca, J, et al; Changes in Myocardial Electrical Impedance in Human Heart Graft Rejection, European Journal of Heart Failure 10 (2008) 594–600
- 21) Wirth, R., et al; Bioelectric Impedance Phase Angle is Associated with Hospital Mortality of Geriatric Patients, Archives of Gerontology and Geriatrics; 2009, 2171, 12.002
- 22) Scanferla, F., et al; Assessment of the Best Compatible Dialysis System: Feasible Application for Bioelectrical Impedance, Int Jnl Art Organs, 1995 Nov; (18)11:712-715
- 23) Doesch, C., et al; Bioimpedance Analysis Parameters and Epicardial Adipose Tissue Assessed by Cardiac Magnetic Resonance Imaging in Patients with Heart Failure, Obesity (2010) doi:10.1038/oby.2010.65
- 24) Oliveira, C., et al; The Phase Angle and Body Cell Mass as Markers of Nutritional Status in Hemodialysis Patients, Journal of Renal Nutrition, March 2010, pp. 1-7

BURDEN OF ILLNESS

- 1) Barbosa-Silva, MCG, et al; Bioelectric Impedance and Individual Characteristics as Prognostic Factors for Post-Operative Complications. Clinical Nutrition (2005) 24, p 830–838
- 2) Zdolsek, HJ, et al; Non-Invasive Assessment of Intercompartmental Fluid s in Burn Victims. Burns 24 (1998) 233-240
- 3) Mueller, PHJ, et al; Predictive Value of APACHE II Score is Improved by Combination with Bioelectrical Impedance Analysis in Multiple Trauma Patients. Critical Care, 1998, 2 (Supplement1): P 162
- 4) Hengstermann, S, et al; Nutrition Status and Pressure Ulcer: What we Need for Nutrition Screening. Journal of Parenteral and Enteral Nutrition, 2007, Jul-Aug; 31(4): p 288-294
- 5) Bottoni, A, et al; Resistance and Reactance in Patients Undergoing Coronary Artery Bypass. Nutricion Hospitalaria, (2003), XVIII (3), p 147-152
- 6) Fritz, T, et al; The Predictive Role of Bioelectrical Impedance Analysis (BIA) in Post-Operative Complications of Cancer Patients. European Journal of Surgical Oncology, 1990; 16: p 326-331
- 7) Gunn, SM, et al; Bioelectrical Phase Angle Values in a Clinical Sample of Ambulatory Rehabilitation Patients. Dynamic Medicine 2008, 7:14
- 8) Mattar, JA, et al; Total Body Impedance Measurements in ARDS. Critical Care Med 1966:A46

- 9) Selberg, O & Selberg, D; Norms and Correlates of Bioimpedance Phase Angle in Healthy Human Subjects, Hospitalized Patients, and Patients with Liver Cirrhosis. European Journal of Applied Physiology 2002; 86:509–516
- 10) Segall, L, et al; Nutritional Status Evaluation and Survival in Hemodialysis Patients in one Centre from Romania. Nephrology, Dialysis and Transplant. 2009 Mar 17. (Electronic publication ahead of print)
- 11) Orea-Tejeda, A, et al; Microalbuminuria in Systolic and Diastolic Chronic Heart Failure Patients. Journal of Cardiology 2008; 15 (2):143-9.
- 12) Barbosa-Silva, MCG, et al; Bioelectrical Impedance Analysis in Clinical Practice: A New Perspective on its Use Beyond Body Composition Equations. Current Opinion in Clinical Nutrition and Metabolic Care 2005, 8:311–317
- 13) Muller, U., et al; Assessment of Body Composition of Patients with COPD. European Journal of Medical Research, 2006 April 28; 11(4):146-51
- 14) Vaninni, FD, et al; Associations Between Nutritional Markers and Inflammation in Hemodialysis Patients. International Urology & Nephrology, DOI 10.1007/s11255-009-9563-8
- 15) Wirth, R., et al; Bioelectrical Impedance Analysis in the Diagnosis of Malnutrition. Journal of Gerontology and Geriatrics, 38:315–321 (2005)
- 16) Marra, M., et al; Bioelectrical Impedance Phase Angle in Constitutionally Lean Females, Ballet Dancers and Patients with Anorexia Nervosa. European Journal of Clinical Nutrition, 2008, November 12.54
- 17) Zamojska, S., et al; Correlates of Habitual Physical Activity in Chronic Haemodialysis Patients. Nephrology, Dialysis and Transplant (2006) 21: 1323–1327
- 18) Bellizzi, V., et al; Early Changes in Bioelectrical Estimates of Body Composition in Chronic Kidney Disease. Journal of the American Society of Nephrology 17: 1481–1487, 2006
- 19) Roos, A.N., et al; Predictive Value of Tetrapolar Impedance Measurements for Hydration for Hydration Status in Critically Ill Patients. Intensive Care Medicine, 1995 February, 21 (2): 125-31
- 20) Goswami, P. N., et al; Bioelectrical Impedance Analysis: Phase Angle - An Independent Predictive Health Marker and its Clinical Applications, Biomed 06, IFMBE Proceedings 15, pp. 321-324, 200.
- 21) Buffa, R, et al; Assessment of Nutritional Status in Free-Living Elderly Individuals Using Bioelectrical Impedance Vector Analysis, Nutrition, 2009, Jan; 25 (1): 3-5.
- 22) Ott M, Fischer H, et al; Bioelectrical Impedance Analysis as a Predictor of Survival in Patients with Human Immunodeficiency Virus Infection. Journal of Acquired Immune Deficiency Syndrome and Human Retrovirology 1995; 9:20–5

- 23) Schwenk A, et al; Phase Angle from Bioelectrical Impedance Analysis Remains an Independent Predictive Marker in HIV-Infected Patients in the Era of Highly Active Antiretroviral Treatment. American Journal of Clinical Nutrition 2000; 72:496–501
- 24) Wagner, D.R., et al; Bioelectrical Impedance as a Discriminator of Pressure Ulcer Risk, Advanced Wound Care, 1996: 9 (2): 30-37.
- 25) Tarulli, A.W., et al; Localized Muscle Impedance Abnormalities in Amyotrophic Lateral Sclerosis, Journal of Clinical Neuromuscular Disease, Volume 10, Number 3, March 2009
- 26) Rutkove, S.B., Electrical Impedance Myography: Background, Current State and Future Directions, Muscle and Nerve, 2009
- 27) Garcia-Lopes, MG, et al; Nutritional Status and Body Composition after 6 Months of Patients Switching from Continuous Ambulatorial Peritoneal Dialysis to Automated Peritoneal Dialysis, Brazilian Journal of Medical and Biological Research (2008) 41: 1116-1122